
Page 1 Weider Nutrition International 2002 South 5070 West Salt Lake City UT 84104

## STRUCTURE / FUNCTION CLAIMS

1300 '98 DEC 18 MO:28

# NUTRIENT: Lactobacillus acidophilus USE THIS REVISED CLAIM REVISED NOVEMBER 5, 1998



DATE: November 5, 1998

BY: Brooke D. Bouwhuis/Luke R. Bucci

Document Name: sflacto05.wpd

BRAND, CODE # & PRODUCT NAME(S):

Schiff® 10500, 10502, 10504 Acidophilus with

Goat Milk (50/100/250 tablets)

**NUTRIENT AMOUNT:** 

Acidophilus Lactobacillus (L. Acidophilus, thermophillus,

bulgaricus) 20,000 CFU; per one tablet

## STRUCTURE/FUNCTION CLAIM:

Digestive Tract Balance

A source of beneficial microorganisms which help maintain a normal balance of healthy flora (bacteria) in the lower intestines for optimal functioning of the digestive tract. Recommended for individuals while taking antibiotics.

### Sales Sheets:

New Schiff® Acidophilus nutritionally supports friendly flora for optimal health and balance in the digestive system. Even the best health care can sometimes fall short of the mark, upsetting the body's natural flora balance. Acidophilus are beneficial microorganisms which help maintain a normal balance of healthy flora (bacteria) for optimal functioning of your digestive system.

- Utilizes Carbohydrates to Survive and Flourish
- Only Carbohydrate in the Woman's Reproductive System, Essential for a Woman's Health
- Fosters Environment Conductive to Helpful Bacteria
- Necessary for Survival in the Small Intestine
- Required for Passage Through the Stomach

Approved by / Date

#### REFERENCES:

- 1. Conway PL, Gorbach SL, Goldin BR. Survival of lactic acid bacteria in the human stomach and adhesion to intestinal cells. J Dairy Sci 1987; 70:1-12.
- Goldin BR, Swenson L, Dwyer J, Sexton M, Gorbach S. Effect of diet and Lactobacillus acidophilus supplements on human fecal bacterial enzymes. J Natl Cancer Instit 1980; 64:255-261.
- Goldin BR, Gorbach SL. Alterations of the intestinal microflora by diet, oral antibiotics, and Lactobacillus: decreased production of free amines from aromatic nitro compounds,

LET2759 62488

Page 2 Weider Nutrition International 2002 South 5070 West Salt Lake City UT 84104

- azo dyes, and glucuronides. J Natl Cancer Inst 1984; 73:689-695.
- 4. Kim GS, Gillialand SE. Lactobacillus acidophilus as a dietary adjunct for milk to aid digestion in humans. *J Dairy Sci* 1983; 66:959-966.
- 5. Kleeman EG, Klaenhammer TR. Adherence of Lactobacillus species to human fetal intestinal cells. *J Dairy Sci* 1982; 65:2063-2069.
- 6. Lilliland SW, Speck ML, Nauyok CF, Giesbrecht FG. Influence of consuming nonfermented milk containing Lactobacillus acidophilus on fecal flora of healthy males. J Dairy Sci 1978; 61:1-10.
- 7. Goldin BR, Gorbach SL. The effect of milk and Lactobacillus acidophilus feeding on human intestinal bacterial enzyme activity. *Am J Clin Nutr* 1984; 39:756-761.
- 8. Friend BA, Shahani KK. Nutritional and therapeutic aspects of lactobacilli *J Appl Nutr* 1984; 36:125-53,
- 9. Guerin-Danan C, Chabanet L, Pedone C, Popot F, Vaissade P, Bouley C, Szylit O, Andrieux C. Milk fermented with yogurt cultures and Lactobacillus casei compared with yogurt and gelled milk: influence on intestinal microflora in healthy infants. *Am J Clin Nutr* 1998; 67:111-7.
- McDonough FE, Hitchins AD, Wong NP, Wells P, Bodwell CE, Modification of sweet acidophilus milk to improve utilization by lactose-intolerant persons. Am J Clin Nutr 1987; 45:570-74.
- 11. Martini MC, Lerebours EC, Lin W, Harlander SK, Berrada NM, Antione JM, Savaino DA. Strains and species of lactic acid bacteria in fermented milks (yogurts): effect on in vivo lactose digestion. *Am J Clin Nutr* 1991; 54:1041–6.